

11/5/21

## Calculus III

## 16.1: Vector Fields

One Last Spherical Example

Ex: Compute the value of <sup>↑ $\theta$</sup> closed disk of radius  $\alpha > 0$ .

NB: We did this already in cartesian coordinates, but it was nasty...

Sol (Spherical Coordinates):  $D_\alpha = \{(p, \theta, \varphi) : 0 \leq p \leq \alpha, 0 \leq \theta \leq 2\pi, 0 \leq \varphi \leq \pi\}$

$$\text{Vol}(D_\alpha) = \iiint_{D_\alpha} 1 \, dV$$

cart

$$= \iiint_{D_\alpha} 1 \cdot p^2 \sin(\varphi) \, dV_{\text{sph.}}$$

spherical

$$= \int_{p=0}^{\alpha} \int_{\theta=0}^{2\pi} \int_{\varphi=0}^{\pi} p^2 \sin(\varphi) \, d\varphi \, d\theta \, dp$$

$$= \int_{p=0}^{\alpha} \int_{\theta=0}^{2\pi} -p^2 [\cos(\varphi)]_{\varphi=0}^{\pi} \, d\theta \, dp$$

$$= \int_{p=0}^{\alpha} \int_{\theta=0}^{2\pi} -p^2 (-1 - 1) \, d\theta \, dp$$

$$= 2 \int_{p=0}^{\alpha} \int_{\theta=0}^{2\pi} p^2 \, d\theta \, dp$$

$$= 2 \int_{\rho=0}^{\alpha} \rho^2 \left[ \theta \right]_{\theta=0}^{2\pi} d\rho$$

$$= 4\pi \int_{\rho=0}^{\alpha} \rho^2 d\rho$$

$$= \frac{4}{3} \pi \left[ \rho^3 \right]_{\rho=0}^{\alpha}$$

$$= \frac{4}{3} \pi (\alpha^3 - 0)$$

$$= \frac{4}{3} \pi \alpha^3$$

## Vector Fields

Goal: Study functions  $\vec{V} : \mathbb{R}^n \rightarrow \mathbb{R}^n$


(most of the time  $n=2$  or  $n=3$  for us)

Defn: A vector field in  $\mathbb{R}^n$  is a function

$$\vec{V} : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Ex: Consider vector field  $\vec{V}(x,y) = \langle x, y \rangle$  on  $\mathbb{R}^2$ . How do we visualize it?


↳ Every point in  $\mathbb{R}^2$  has an associated vector attached to it by this vector field:



\* draw the vector  $\vec{V}(x,y)$  with its tail at point  $(x,y)$  \*

$$\vec{V}(1,0) = \langle 1, 0 \rangle$$

Ex: Draw  $\vec{v}(x,y) = \langle -y, x \rangle$ .



$$\begin{aligned} v(1,0) &= \langle 0, 1 \rangle \\ v(2,0) &= \langle 0, 2 \rangle \\ v(-1,0) &= \langle 0, -1 \rangle \\ v(-2,0) &= \langle 0, -2 \rangle \\ v(-1,1) &= \langle -1, -1 \rangle \\ v(0,1) &= \langle -1, 0 \rangle \\ v(-2,1) &= \langle -1, -2 \rangle \end{aligned}$$

Ex: The gradient of a function is always a vector field.

vector field = "v.f."

↳ e.g. for  $f(x,y) = xy$ ,  $\nabla f = \langle y, x \rangle$  is a vector field on  $\mathbb{R}^2$

- A vector field like this is sometimes called a "gradient vector field".

e.g.  $f(x,y,z) = e^{x+y^2} \cos(z+x)$

$$\nabla f = \langle e^{x+y^2} \cos(z+x) - e^{x+y^2} \sin(z+x), 2ye^{x+y^2} \cos(z+x), -e^{x+y^2} \sin(z+x) \rangle$$

is a vector field  $\vec{\nabla} f(x,y,z)$

• Obvious Question: How do we know when a vector field is a gradient vector field?

↳ Is every v.f. a grad. v.f.?

Terminology: A conservative vector field is a gradient v.f.

If  $\vec{V} = \nabla f$  is a conservative v.f., we call  $f$  a potential function for  $\vec{V}$

Now, is every v.f. conservative?

In  $\mathbb{R}^2$ , a conservative v.f. has form

$\vec{v} = \langle f_x(x, y), f_y(x, y) \rangle$  for some potential function  $f$ . By Clairaut's Theorem,  $f_{xy} = f_{yx}$ , so every conservative v.f.  $\vec{v} = \langle \alpha(x, y), \beta(x, y) \rangle$  has to satisfy  $\alpha_y = \beta_x$ .

Ex:  $\vec{v} = \langle \overset{v_x}{-y}, \overset{v_y}{x} \rangle$  has

$$\frac{d}{dy} [v_x] = \frac{d}{dy} [-y] = -1$$

On the other hand ...

$$\frac{d}{dx} [v_y] = \frac{d}{dx} [x] = 1$$

$\therefore \vec{v}$  is non conservative lest it violates Clairaut's Theorem

$\therefore$  Not every vector field is a gradient vector field.  $\cap$

Prop: A vector field  $\vec{v} = \langle v_{x_1}, v_{x_2}, \dots, v_{x_n} \rangle$  is conservative if and only if  $\frac{d}{dx_i} [v_{x_j}] = \frac{d}{dx_j} [v_{x_i}]$

for all  $i, j$ . (i.e. a v.f. is conservative if and only if it satisfies Clairaut's Theorem)

Ex: Is  $\vec{v} = \langle x, y \rangle$  conservative?

Sol:  $\frac{dv_x}{dy} = \frac{d}{dy} [x] = 0$

$$\frac{dv_y}{dx} = \frac{d}{dx} [y] = 0$$

∴ by the proposition  $\vec{V} = \langle x, y \rangle$  is conservative:

What is its potential function?

By conservativity,  $\nabla f = \vec{V}$  for some function  $f(x, y)$ . i.e.  $f_x(x, y) = x$ , and  $f_y(x, y) = y$ .

$$\text{B/c } \frac{df}{dx} = x, \text{ we know } f(x, y) = \int \frac{df}{dx} dx = \int x dx \\ = \frac{1}{2}x^2 + c(y)$$

$$\therefore y = \frac{df}{dy} = \frac{d}{dy} \left[ \frac{1}{2}x^2 + c(y) \right] = \frac{df}{dy}$$

$$\therefore c(y) = \int \frac{dc}{dy} dy = \int y dy = \frac{1}{2}y^2 + D$$

← An actual constant (honest constant)

$$\text{Hence } f(x, y) = \frac{1}{2}x^2 + c(y) = \frac{1}{2}x^2 + \frac{1}{2}y^2 + D$$

← for any constant

$D$  is a potential function of  $\vec{V}$

Ex:  $\vec{V} = \langle 2xy, x^2 - 3y^2 \rangle$ . Conservative? If yes, potential?

$$\text{Sol: } \frac{dV_y}{dx} = \frac{d}{dx} [x^2 - 3y^2] = 2x$$

$$\frac{dV_x}{dy} = \frac{d}{dy} [2xy] = 2x$$

∴  $\vec{V}$  is conservative

If  $\vec{V} = \nabla f$ , then  $f_x = 2xy$

$$f = \int f_x dx = \int 2xy dx = x^2 y + c(y)$$

$$\therefore f_y = \frac{d}{dy} [x^2 y + c(y)]$$

$$x^2 - 3y^2 = x^2 + c'(y)$$



$$\therefore c'(y) = -3y^2$$

$$\therefore c(y) = \int -3y^2 dy = -y^3 + D$$

$$\therefore f(x,y) = x^2y + c(y) = x^2y - y^3 + D$$

← is a potential for all constant D

NB: The method for computing the potentials can be used to prove the proposition from earlier...

Ex:  $\vec{v} = \langle \ln(x+y), e^{x+y} + \frac{1}{x+y} \rangle$

Conservative? If yes, potential?

Sol:  $\frac{dv_x}{dy} = \frac{d}{dy} [\ln(x+y)] = \frac{1}{x+y}$

$$\frac{dv_y}{dx} = \frac{d}{dx} \left[ e^{x+y} + \frac{1}{x+y} \right] = e^{x+y} - (x+y)^{-2}$$

$\therefore \vec{v}$  is not conservative,  $\therefore$  no potential.

Last Time: You saw

$$\int_{y=c}^d \int_{x=a}^b f(x)g(y) dx dy = \int_{x=a}^b f(x) dx \cdot \int_{y=c}^d g(y) dy$$

only true when:

- ① Integrating over a rectangle  $[a,b] \times [c,d]$
- ② Integrated as a "separable function" i.e.  $h(x,y) = f(x) \cdot g(y)$